



Assessment of Heavy Metals (Fe, Cu and Ni)  
Contamination of Seawater and Mussel, *Mytilus*  
*Galloprovincialis*, from Al Hoceima Moroccan  
Coasts

---

Ghizlane Azizi, Mostafa Layachi, Mustapha Akodad,  
Hamza Ngadi, Mourad Baghour, Ali Skalli, Mohammad Ghalit,  
Elkhadir Gharibi and Abdelmajid Moumen

EasyChair preprints are intended for rapid  
dissemination of research results and are  
integrated with the rest of EasyChair.

March 3, 2020

# Assessment of heavy metals (Fe, Cu and Ni) contamination of seawater and mussel, *Mytilus galloprovincialis*, from Al Hoceima Moroccan coasts

Heavy metal concentrations in *Mytilus galloprovincialis*

Ghizlane Azizi<sup>\*1</sup>, Mostafa Layachi<sup>2</sup>, Mustapha Akodad<sup>1</sup>, Hamza Ngadi<sup>1</sup>, Mourad Baghour<sup>1</sup>, Ali Skalli<sup>1</sup>, Mohammad Ghalit<sup>3</sup>, Elkhadir Gharibi<sup>3</sup>, Abdelmajid Moumen<sup>1</sup>

<sup>1</sup>OLMAN- RL, Faculté Pluridisciplinaire Nador (FPN), Nador, Université Mohammed 1<sup>er</sup>, Maroc.

<sup>2</sup>Centre Régional de l'INRH-Nador-, 13 Boulevard Zerktouni BP: 493, Nador, Maroc.

<sup>3</sup>LCSMA, Faculté des Sciences Oujda, Université Mohammed 1<sup>er</sup>, Maroc.

\*Corresponding authors: e-mail: [ghizlaneazizi@hotmail.com](mailto:ghizlaneazizi@hotmail.com)

## ABSTRACT

In the present work, both seawater and mussels (*M. galloprovincialis*) were collected monthly from the mussel farm located along the coastline of Al Hoceima (Morocco) and their heavy metal concentrations (Fe, Cu, Ni) were determined using inductively coupled plasma – optical emission spectrometry (ICP-OES). The trace metals found in the mussel tissues are much higher than those found in the environment. The order of the abundance of metals in the mussels is Fe > Cu > Ni. The pattern of the temporal variation of mussel trace metal levels was similar among stations with higher values during the months of winter season and lower during the months of summer period. Heavy metals uptake is dependent on both environmental and biological factors. Results obtained in this study for Fe, Cu and Ni, indicate that the species under investigation poses no health hazard to consumers because its trace metal contents remain within the permissible range established by various international guideline for safe human consumption. Our study, indicate that bivalve molluscs can be used as sentinel bioindicators in marine environments impacted by heavy metals in this area of the Moroccan coastline.

**KEYWORDS** Heavy metals, seawater, Moroccan Mediterranean coastline, *Mytilus galloprovincialis*, humain health.

## INTRODUCTION


The rapidly increasing population and developing industrial and agricultural activities in mediterranean countries have caused increase in industrial, agricultural and municipal water pollution. The discharge of effluent, treated or not treated, into the sea and rivers has resulted in soil and water pollution. The anthropogenic activities wastewater contains not only organic matter, but also other pollutants such as heavy metals [1]. These trace elements occur in the marine environment to levels that are hazardous for marine biota and can be dangerous for human health by consuming these marine organisms. A number of national and international programs have been established over recent decades to identify efficient and accurate biomonitors of trace metal pollutant availabilities [2, 3]. Extensive surveys have been undertaken in many countries [4, 5]. Authors (Pérez-López et al. [6]; Yuan et al. [7]) reviewed the use of biological indicators to monitor chemical pollutants such as trace metals and concluded that such organisms may have advantages over classical methods of water and sediment analysis. Mussels are widely used as sentinel organisms in marine pollution monitoring programs. In this occurrence, mussels have been proposed as biomonitors because of their wide geographical distribution, easy sampling, sedentary nature, filtering habits, resistance to stress, sessile lifestyle and ability to bioaccumulate heavy metals in their tissues to levels many times higher than their surrounding water [8-10]. In the bivalve molluscs, trace metals have been absorbed both from water and from ingested phytoplankton and other suspended particles [2]. Thus, trace metals accumulation in these marine organisms can be affected by environmental (salinity, temperature, intertidal zone, available food, etc) and physiological (age, reproductive cycle, sex, etc.) factors [11, 12]. No research, however, has been carried out so far on seasonal variation in metal accumulation or on biological factors affecting metal accumulation in mussel populations from the mediterranean moroccan coasts.

The goal of this study was to determine the monthly levels of Fe, Cu and Ni in the soft tissue of mussels *Mytilus galloprovincialis* and seawater samples along the Al Hoceima coastline. The results of this study may provide information on the use *Mytilus galloprovincialis* as biological indicators of heavy metals in this part of the Mediterranean coasts. The work studies also the levels of toxic metals in the mussels and if they are within the maximum permissible levels set by the international guidelines.

## MATERIAL AND METHODS

### 1. Area of study, collection of samples and preparation of mussels

The Mediterranean mussel *M. galloprovincialis* and seawater were obtained from the mussel farming installed in the coastal areas of Al Hoceima. Waters and mussel samples were collected monthly in 2018, from January to December, from five sampling sites (A, B, F, J and I) (Fig. 1), to ensure homogeneous distribution of samples. Water was sampled by hand into several bottles according to the analytical specifications and transported at +4°C in a cold box to the laboratory until analysis.



**Figure 1: Geographical location of mussel sampling sites along the mussel farming from Al Hoceima coastline (Moroccan Mediterranean).**

Approximate 50 mussels (*Mytilus galloprovincialis*) from the sampling sites were collected. The mean value  $\pm$  standard deviation (SD) of the total width, length and whole weight of samples was  $39.3 \pm 0.6$  mm,  $66.1 \pm 1.1$  mm and  $29.1 \pm 1.3$  g, respectively. The specimens were immediately washed at each site with seawater to eliminate encrusted organisms, stored in bags, kept in a cooler box with ice and transported to the laboratory. For a 48-hour period, the mussels were kept alive in clean seawater to purge their digestive systems, in order to be able to measure only the heavy metals biologically deposited in their tissues [13, 12].

The whole soft tissues of mussels were removed from the shells using a steel knife, then thoroughly rinsed with distilled water to remove the extraneous impurities. Whole tissues were dried at 60 °C for 48 h to obtain the dry weight, and were ground in a mortar for homogenization. The ground powder of mussel tissues were grouped in five pools monthly (each pool corresponds to five individuals of similar size from each sites) for chemical analyses of metals.

## 2. Metals analysis

The metal contents (Fe, Cu and Ni) in the whole soft tissues of mussels and water samples were analyzed by inductively coupled plasma – optical emission spectrometry (ICP-OES 720-ES), (Agilent Technologies, USA) in the Scientific Instrumentation Service at the Estación Experimental del Zaidín, Consejo Superior de Investigaciones Científicas (EEZ-CSIC), Granada, Spain.


Approximately 0.3 g of dry weight of the mussel soft tissues was digested by Start D Microwave Digestion System (Milestone), equipped with a rotor MPR SK-12T (Milestone), using a 10 ml of solution of 1:4 (H<sub>2</sub>O<sub>2</sub> (30%) : HNO<sub>3</sub> (65%)), at 190 °C, during 1 h 30 min as described by Belivermiş et al. [14]. An external calibration was performed for quantification, which utilizes different multielement solutions to cover the targeted analyte range [2].

## 3. Statistical analysis

The data were expressed as means and standard deviations, and treated using one-way analysis of variance (ANOVA), followed by Tukey's test, accepting  $P < 0.05$  classified as statistically significant.

# RESULTS AND DISCUSSION

## 1. Trace metals in the seawater of Al Hoceima coastline



**Figure 2: Monthly variations of trace metals (Fe, Cu and Ni) in seawater from Al Hoceima coastline (Moroccan Mediterranean), during the sampling period (displayed as mean  $\pm$  standard deviation).**

**Table 1. Comparison of heavy metal concentrations as mg/kg in seawater from Al Hoceima coastline with the permissible limits recommended by the FAO.**


	Cu	Fe	Ni
Present Study	0.0905	0.3545	0.1018
FAO [15]	0.5	---	0.5
FAO/WHO [16]		5.0	

The average monthly trace metals of seawater from Al Hoceima coastline areas, during the sampling period of 2018 are shown in Figure 2. The pattern of accumulation in seawater was Fe > Ni > Cu. Iron (Fe) in sea water present a peak in December ( $1.2930 \pm 0.018$  mg/kg). The lowest value of Ni content was also observed in April ( $0.0037 \pm 0.0001$  mg/kg) and the maximum values during September ( $0.1869 \pm 0.0186$  mg/kg). The highest Cu concentration was recorded in autumn with a peak in November ( $0.1579 \pm 0.011$  mg/kg), and the minimum value was observed in January ( $0.0411 \pm 0.0029$  mg/kg). Trace metal concentrations in seawater of Al Hoceima coastal regions were quite similar to values referred to in other regions of the world coastal waters [17, 6]. The pattern may be the result of local anthropogenic activities, due to industrial, agricultural and domestic sources. In this study, trace metal levels were found to be lower than the recommended limits

established by various international guidelines (Table 1) [16, 15]. These measurements are important to show the metal concentrations in the water and to make a correlation with those in the mussels.

The heavy metals found in the mussel tissues (Fig. 3, see below) are much higher than those found in their marine environment. Similar results are found in previous studies [6, 18]. Mussels absorb trace metals both from water and from ingested phytoplankton and other suspended particles [2, 19]. Because of their tendency to concentrate different pollutants from their environment, marine bivalves have shown useful as sentinel organisms, since the levels of heavy metal contaminations in the molluscs are directly proportional to the level of their availability in the environment [6, 20].

## 2. Seasonal trends of Zn, Cd and Cr in soft tissues of *Mytilus galloprovincialis*



**Figure 3: Monthly changes in soft tissue metal (Fe, Cu and Ni) contents (mg/kg dry weight) in *Mytilus galloprovincialis* from Al Hoceima coastline (Morocco), (mean  $\pm$  standard deviation).**

Monthly mean concentrations of each metal in the soft tissue of *M. galloprovincialis* are presented graphically (Fig. 3). Taking into account the highest concentration of metals in any of the months, the orders of decreasing concentration in soft tissues of mussels were Fe > Cu > Ni.

Average metal (Fe, Cu and Ni) concentrations in the whole soft tissues of mussels were ( $542.57 \pm 47.15$ ,  $7.193 \pm 0.142$  and  $3.502 \pm 0.251$  mg/kg, for Fe, Cu and Ni; respectively).

The Fe levels in soft tissues of mussels present higher values in winter period with a peak in February ( $1022.83 \pm 6.750$  mg/kg), the minimum and similar values were observed in summer-autumn seasons. The highest Cu content over the period analyzed occurred in February ( $6.955 \pm 0.234$  mg/kg) and the lowest in August ( $4.455 \pm 0.370$  mg/kg). Elevated concentrations of Ni occurred in winter period with a peak in December ( $4.545 \pm 0.132$  mg/kg) and the lowest value was observed in summer season for the month of August ( $1.457 \pm 0.175$  mg/kg).

In the present study, the seasonal fluctuations of soft tissue metal concentrations in mussels may be driven by various environmental and biological factors. Changes in trace metal contents (Fe, Cu and Ni) found in *M. galloprovincialis*, with higher values in mussels collected in the winter and lower contents were observed in soft tissues of mussels in the summer, have long been considered [21, 5, 22, 23]. The influence of seasonal changes in metal bioaccumulation in mussels can be explained in terms of the characteristics of climate change in the Mediterranean area (warm, dry in summers and wet, rainy in winters). Of the processes affecting metal dynamics in mussels, variability in physicochemical conditions of the water and sediments. The period of winter is known as a rainy season. During this period, rivers collect water through branched drainage canals from large inland areas and discharge it to the sea, often with great quantities of terrestrial inputs from the studied region. The coastal marine environment of Al Hoceima region receive a large amount of suspended matter and systematically large charge of particle-associated elements of various kinds including heavy metals, that can originate from natural sources (processes of chemical weathering and geochemical activity) and from local pollution sources (agriculture, municipal wastewater and industrial activity). In addition, the increase of metal (Fe, Cu and Ni) concentrations in mussel tissues during the winter period, may have been associated with faster metal uptake and accumulation during intense food uptake of the mussels in this period [24]. Trophic transfer of metals is increasingly recognized as an important pathway for metal accumulation in marine bivalves [25]. In Other studies, the seasonal changes in trace metal contents in marine biota have been exclusively attributed to variations in the levels of these elements in water [3, 19].

Another factor which affect the metal contents in mussels is the salinity gradient. Authors (Blust et al. [26]; Blackmore and Wang [27]) indicated that trace metal bioavailability decreases with increasing salinity, according to the concentrations of free metal ion species in the exposure water. Authors (Azizi et al. [19]; Cinnirella et al. [28]) found that in the Mediterranean coasts, the concentration of traces metals increase and reach its maximum in the winter when the salinity levels decrease and reach its lowest level.

The seasonal trends of trace metal concentrations in mussels can also be explained in terms of changes in physiological status (reproductive stage, sex, age, growth, etc). Among physiological processes, reproduction is one of the most energy demanding and can result in energetic trade-offs and modulation of physiological performance [5]. Phillips [29] reported that seasonal fluctuations in heavy metal concentrations in mussel are mainly due to dry weight fluctuations against a background of fairly stable metal content throughout a year. The effect of weight changes on metal levels has been also established by authors (Sokolowski et al. [30]) in marine bivalves. Furthermore, many authors (Widdows [31]; Dokmeci, [12]) have suggested that tissue metal concentrations in mussels vary considerably according to reproductive stage. Azizi et al. [19] found that up to 40% of tissues were lost during the spawning period.

**Table 2. Comparison of heavy metal concentrations as  $\mu\text{g}/\text{kg}$  dry weight in mussels, *Mytilus galloprovincialis*, from the diverse coastal locations with the tolerance levels.**

Location	Cu	Ni	Fe	Références
Cala Iris Sea	7.193	3.502	542.57	Présent work
Saronikos gulf, Greece	20.0	4.9	593	Strogloudi et al. [5]
Western Scheldt estuary (Netherlands)	27.4	16.2	572	Mubiana et al. [32]
Chacopata- Bocaripo Lagoon (Venezuela)	9.00	8.47		Pinto et al. [33]
Apulian Coast (Italy)	8.17			Spada et al. [2]
ElJadida, Moroccan coast	142.2	94.3		Maanan [34]
Galicia and Cantabria, Spain	10.09			Besada et al. [11]
Brown Bay, Argentina	2.00		144.69	Giarratano and Amin [21]
Island of Sylt, Germany	8.272	2.007		Helmholz et al. [35]
Baltic Sea (norway)	12.10	9.60	1130	Pempkowiak et al. [36]
Norwegian Sea	7.57	3.49	475	Pempkowiak et al. [36]
Moroccan Atlantic coast	43.1			Maanan [13]
Andalucia coast (Spain)	11.9			Benedicto et al. [37]
Galicia and Gulf of Biscay (NW Spain)	14.06			Besada et al. [38]
Croatian Coasts (Adriatic Sea)	10.53			Kljaković-Gaspić et al. [39]
Montenegro Coasts (Adriatic Sea)	8.26			Joksimovic et al. [40]
Baleari Islands (Western Mediterranean)	4.76			Deudero et al. [41]
Guideline	50-150			USEPA [42]
Guideline	70			FAO [43]
Guideline		80		USFDA [44]
Guideline	20			TKB [45]
Guideline	20			UNEP [46]

The comparison of the some heavy metal concentrations obtained in the mussel samples collected from the Al Hoceima coastal region with the data from different areas and with the tolerance levels in mussels given by the international standards was given in Table 2.

The trace metal concentrations in mussels, expressed in mg/kg dry weight, for Fe, Cu, and Ni are: 542.57, 7.193 and 3.502, respectively. Copper is an essential mineral for human health and at the same time can be toxic, depending upon the amounts ingested [47]. In this sense, a severe deficiency of copper has been associated with bone malformation during development [48], and as a contributory factor to osteoporosis in adults [49], it has also been associated with cardiovascular risk and altered immune responses [50]. However, high uptakes of copper may cause liver and kidney damage and even death [51]. Nickel is a hard metal found in soil as well as in foods. In small quantities nickel is essential, but when the uptake is too high it can be a danger to human health. This compound can cause a variety of adverse effects on human health such as dermatitis, chronic bronchitis, cardiovascular and kidney diseases, and cancer of the lung and nasal sinus [52]. Iron is also an essential element for almost all living organisms as it participates in a wide variety of metabolic processes, including oxygen transport, deoxyribonucleic acid (DNA) synthesis, and electron transport. However, in excess quantities, Iron can also be poisonous and cause serious threats to human health such as conjunctivitis, choroiditis, retinitis, and also participate in development of a siderosis [53]. Based on a comparison of the permissible limits set by various international organizations, Cu levels in this study were lower than the Food and Agriculture Organization permissible limit which is 70 mg/kg [43], and the United Nations Environment Programme [46] and the Turkish [45] permissible limit which is 20 mg/kg. On the other hand, according to the United States Environmental Protection Agency [42], the maximum permissible Cu levels are 50-150 mg/kg for molluscs. The Ni contents in the present study were lower than the United States Food and Drug Administration permissible limit which is 80 mg/kg [44]. However, for Fe, there is no information about maximum permissible iron concentrations in fish/molluscs tissues recommended by various organizations (Table 2).

In general, metal levels of the present study were in agreement with the results of previous mussel studies in diverse coastal locations. Copper (Cu) levels obtained in this study were lower than those reported for ElJadida, Moroccan coast [34], Western Scheldt estuary of Netherlands [32], Saronikos gulf of Greece [5], Galicia and Gulf of Biscay of Spain [38] and Croatian Coasts of Adriatic Sea [39], but higher than those reported for Baleari Islands of Western Mediterranean [41] and for Brown Bay of Argentina [21]. On the other hand, our findings for Cu levels were in agreement with ones reported for Norwegian Sea [36]. For Ni, the concentrations found in *M. galloprovincialis* in the present work, were in the range of the literature which are 2.01-94.3 mg/kg, in tissues of *Mytilus galloprovincialis* from Island of Sylt, Germany [35] and El Jadida, Moroccan coast [34]; respectively (Table 2). Iron (Fe) levels found in tissues of mussels from Cala Iris Sea, were in the range of the bibliographic works, which are 144.69-1130 mg/kg, in soft tissues of mussel from Brown Bay of Argentina [21] and Baltic Sea of Norway [36]; respectively (Table 2).

## CONCLUSION

Our studies reveal interesting results concerning the bioaccumulation of Fe, Cu and Ni in *Mytilus galloprovincialis* from the Moroccan Mediterranean coasts during the season period. Trace metal levels in soft tissues of mussels present higher values in winter period because of the Mediterranean climate change, river flow discharges chemical pollutant from local pollution sources (industrial, agricultural and domestic activities), and also of the physiological status of the bivalve (age, weight and reproductive cycle). The results obtained in this study were compared with those reported in earlier studies and concluded that, the mussel from Al Hoceima coastline was in general not considered a metal polluted mollusc according to the international guidelines (FAO, USFDA, TKB, USEPA, UNEP). So, the Fe, Cu and Ni, under investigation in this work, pose no health hazard to consumers. The heavy metals found in soft tissues of mussel are much higher than those found in their marine habitats. We concluded that the marine biota can be used as bioindicator for monitoring of coastal water pollution in the region of Morocco.

## REFERENCES

- [1] Zhang, Y., Wang, M., Huang, B., Akhtar, M.S., Hu, W. and Xie, E., 2018. Soil mercury accumulation, spatial distribution and its source identification in an industrial area of the Yangtze Delta, China. *Ecotoxicol Environ Saf*, 163:230–237 <https://doi.org/10.1016/j.ecoenv.2018.07.055>.
- [2] Spada, L., Annicchiarico, C., Cardellicchio, N., Giandomenico, S. and Di Leo, A., 2013. Heavy metals monitoring in the mussel *Mytilus galloprovincialis* from the Apulian coast (Southern Italy). *Mediterr Mar Sci*, 14:99-108.
- [3] Lacroix, C., Duvieilbourg, E., Guillou, N., Guyomarch, J., Bassoulet, C., Moraga, D., Chapalain, G. and Auffret, M., 2017. Seasonal monitoring of blue mussel (*Mytilus* spp.) populations in a harbor area: A focus on responses to environmental factors and chronic contamination. *Mar Environ Res*, 129: 24-35.
- [4] Szefer, P., Geldon, J., Ali, A.A., Bawazir, A. and Saad, M., 1997. Distribution and association of trace metals in soft tissue and byssus of mollusc *Perna perna* from the Gulf of Aden, Yemen. *Environ Int*, 2:53–61.
- [5] Stroglyoudi, E., Angelidis, M.O., Christides, A. and Papatthassiou, E., 2012. Metal concentrations and metallothionein levels in *Mytilus galloprovincialis* from Elefsis bay (Saronikos gulf, Greece). *Environ Monit Assess*, 184:7189–7205.
- [6] Pérez-López, M., Alonso, J., Nóvoa-Valiñas, M.C. and Melgar, M.J., 2003. Assessment of heavy metal contamination of seawater and marine limpet, *Patella vulgata* L., from Northwest Spain. *J Environ Sci Health A Tox Hazard Subst Environ Eng*, 12:2845–2856.
- [7] Yuan, Y., Sun, T., Huijuan, W., Liu, Y., Pan, Y., Xie, Y., Huang, H. and Fan, Z., 2019. Bioaccumulation and health risk assessment of heavy metals of bivalve species in Daya Bay (South China Sea): Consumption advisory. *Mar Pollut Bull*, Doi: MPB-D-1901012.
- [8] Goldberg, E.D., 1975. The mussel watch: a first step in global marine monitoring. *Mar Pollut Bull*, 6:111–132.
- [9] Fernandez-Tajes, J., Florez, F., Pereira, S., Rabade, T., Laffon, B. and Mendez, J., 2011. Use of 3 bivalve species for biomonitoring a polluted estuarine environment. *Environ Monit Assess*, 177:289–300.
- [10] Dokmeci, A.H., Yildiz, T., Ongen, A. and Sivri, N., 2014. Heavy metal concentration in deep water rose shrimp species (*Parapenaeus longirostris*, Lucas, 1846) collected from the Marmara Sea Coast in Tekirdağ. *Environ Monit Assess*, 186:2449–2454.
- [11] Besada, V., Andrade, J.M., Schultze, F. and Gonzalez, J.J., 2011. Monitoring of heavy metals in wild mussels (*Mytilus galloprovincialis*) from the Spanish North-Atlantic coast. *Cont Shelf Res*, 31:457–65.
- [12] Dokmeci, A.H., 2017. Assessment of heavy metals in wild mussels *Mytilus galloprovincialis* from the Marmara Sea Coast of Tekirdag (Turkey). 15th International Conference on Environmental Science and Technology Rhodes, Greece, 31 August to 2 September 2017. CEST2017\_00569.
- [13] Maanan, M., 2007. Biomonitoring of heavy metals using *Mytilus galloprovincialis* in Safi coastal waters Morocco. *Environ Toxicol*, 22:525–531.
- [14] Belivermiş, M., Kılıç, O. and Çotuk, Y., 2016. Assessment of metal concentrations in indigenous and caged mussels (*Mytilus galloprovincialis*) on entire Turkish coastline. *Chemosphere*, 144:1980–1987.
- [15] FAO, 2006. Elaboration des fiches techniques des valeurs limites des rejets industriels. Projet de gestion des ressources en eau: Elaboration des dossiers techniques relatifs aux valeurs limites des rejets industriels dans le Domaine Public Hydraulique. Convention FAO/UTF/MOR019/MOR, 1-23.
- [16] FAO/WHO, 1985. Joint/WHO Expert Committee on Food Additives 1999. Summary and Conclusions. Proceedings of the 53rd Meeting, Rome, June, 1-10.
- [17] Paek, S.M., Chung, S. and Lee, I.S. 1999. Level of heavy metals in the Oasan Bay in Korea and involvement of metal binding protein in the accumulation of cadmium in *Littorina brevicula*. *Korean J Ecol*, 22(2):95–100.

- [18] Bilandžić, N., Sedak, M., Čalopek, B., Džafić, N., Ostojić, D.M. and Potočnjak, D. 2015. Metal content in four shellfish species from the Istrian coast of Croatia. *Bull Environ Contam Toxicol*, 95:611–617.
- [19] Azizi, G., Akodad, M., Baghour, M., Layachi, M. and Moumen, A., 2018. The use of *Mytilus* spp. mussels as bioindicators of heavy metal pollution in the coastal environment. A review. *J Mater Environ Sci*, 9 1170-1181.
- [20] Rainbow, P.S., 2007. Trace metal bioaccumulation: models, metabolic availability and toxicity. *Environ Int*, 33:576-582.
- [21] Giarratano, E. and Amin, O.A., 2010. Heavy metals monitoring in the southernmost mussel farm of the world (Beagle Channel, Argentina). *Ecotox Environ Safe*, 73(6):1378–1384.
- [22] Shue, M.-F., Chen, W.-D., Bellotindos, L.-M. and Lu, M.-C., 2014. Seasonal variations of heavy metals content in muscle and viscera of green-lipped mussel *Perna viridis* from Da-Peng Bay lagoon in Taiwan. *J Toxicol Environ Health A*, 77:1222–1228.
- [23] Piscopo, M., 2019. Seasonal dependence of cadmium molecular effects on *Mytilus galloprovincialis* (Lamarck, 1819) protamine- like protein properties. *Mol Reprod Dev*, 1–12. doi: 10.1002/mrd.23240.
- [24] Bocchetti, R. and Regoli, F., 2006. Seasonal variability of oxidative biomarkers, lysosomal parameters, metallothioneins and peroxisomal enzymes in the mediterranean mussel *Mytilus galloprovincialis* from Adriatic Sea. *Chemosphere*, 65(6):913–921.
- [25] Scudiero, R., Creti, P., Trinchella, F. and Esposito, M.G., 2014. Evaluation of cadmium, lead and metallothionein contents in the tissues of mussels (*Mytilus galloprovincialis*) from the Campania coast (Italy): Levels and seasonal trends. *C R Biol*, 337:451-458.
- [26] Blust, R., Kockelbergh, E. and Baillieul, M., 1992. Effect of salinity on the uptake of cadmium by the brine shrimp *Artemia franciscana*. *Mar Ecol Prog Ser*, 84:245–254.
- [27] Blackmore, G. and Wang, W.-X., 2003. Inter-population differences in Cd, Cr, Se, and Zn accumulation by the green mussel *Perna viridis* acclimated at different salinities. *Aquat Toxicol*, 62:205–218.
- [28] Cinnirella, S., Bruno, D.E., Pirrone, N., Horvat, M., Živković, I., Evers, D.C., Johnson, S. and Sunderland, E.M., 2019. Mercury concentrations in biota in the Mediterranean Sea, a compilation of 40 years of surveys. *Sci Data* 6:205 <https://doi.org/10.1038/s41597-019-0219-y>.
- [29] Phillips, D.J.H., 1980. Quantitative aquatic biological indicators: Their use to monitor trace metal and organochlorine pollution. Applied Science Publishers, London.
- [30] Sokolowski, A., Bawazir, A.S. and Wolowicz, M., 2004. Trace metals in the brown mussel *Perna perna* from the coastal waters off Yemen (Gulf of Aden): how concentrations are affected by weight, sex, and seasonal cycle. *Arch Environ Contam Toxicol*, 46:67–80.
- [31] Widdows, J., 1985. Physiological responses to pollution. *Mar Pollut Bull*, 16:129-134.
- [32] Mubiana, V.K., Qadah, D., Meys, J. and Blust, R., 2005. Temporal and spatial trends in heavy metal concentrations in the marine mussel *Mytilus edulis* from the Western Scheldt estuary (The Netherlands). *Hydrobiologia*, 540:169–180.
- [33] Pinto, R., Acosta, V., Segnini, M.I., Brito, L. and Martínez, G., 2015. Temporal variations of heavy metals levels in *Perna viridis*, on the Chacopata-Bocaripo lagoon axis, Sucre State, Venezuela. *Mar Pollut Bull*, 91:418–423.
- [34] Maanan, M., 2008. Heavy metal concentrations in marine molluscs from the Moroccan coastal region. *Environ Pollut*, 153:176-183.
- [35] Helmholtz, H., Ruhnau, Ch., Pröfrock, D., Erbslöh, H.B. and Prange, A., 2016. Seasonal and annual variations in physiological and biochemical responses from transplanted marine bioindicator species *Mytilus* spp. during a long term field exposure experiment. *Sci Total Environ*, 565:626–636.
- [36] Pempkowiak, J., Sikora, A. and Biernacka, E., 1999. Speciation of heavy metals in marine sediments vs their bioaccumulation by mussels. *Chemosphere*, 39:313-321.
- [37] Benedicto, J., Rodriguez, C., Martinez-Gomez, C., Guerrero, J. and Jornet, A., 2003. Distribucion espacial y tendencias temporales de los niveles de metales traza en el litoral de Andalucía utilizando mejillon *Mytilus galloprovincialis* Lamarck, 1819 como organismo indicador: 1991–2003. *Bol Inst Esp Oceanogr*, 19:31–39.
- [38] Besada, V., Andrade, J.M., Schultze, F., Fumega, J., Cambeiro, B. and Gonzalez, J.J., 2008. Statistical comparison of trace metal concentrations in wild mussels (*Mytilus galloprovincialis*) in selected sites of Galicia and Gulf of Biscay (Spain). *J Mar Syst*, 72:320–331.
- [39] Kljaković-Gaspić, Z., Herceg-Romanić, S., Kozul, D. and Veza, J., 2010. Biomonitoring of organochlorine compounds and trace metals along the Eastern Adriatic coast (Croatia) using *Mytilus galloprovincialis*. *Mar Pollut Bull*, 60:1879-1889.
- [40] Joksimovic, D., Tomic, I., Stanković, A.R., Jovic, M. and Stanković, S., 2011. Trace metal concentrations in Mediterranean blue mussel and surface sediments and evaluation of the mussels quality and possible risks of high human consumption. *Food Chem*, 127:632-637.
- [41] Deudero, S., Box, A., Tejada, S. and Tintoré, J., 2009. Stable isotopes and metal contamination in caged marine mussel *Mytilus galloprovincialis*. *Mar Pollut Bull*, 58:1025-1031.
- [42] United States Environmental Protection Agency (USEPA), 2002. Appendix V: Median International Standards. Environmental Protection Agency. State Water Resources Control Board. <http://www.swcb.ca.gov>.
- [43] FAO, 1983. Compilation of legal limits for hazardous substances in fish and fishery products, FAO fishery circular, No. 464, pp. 5–100.
- [44] United States Food And Drug Administration (USFDA), 1993. Guideline for toxic elements. Chapter 23: Environmental chemical contaminants and pesticides. In Seafood Network Information Center, Sea Grant Extension Program. Sea Grant, California.
- [45] TKB, 2002. Fisheries laws and regulations. Ministry of Agriculture and Rural Affairs, Conservation and Control General Management, Ankara.
- [46] United Nations Environment Programme (UNEP), 1993. Guidelines for monitoring chemical contaminants in the sea using marine organisms. UNEP/FAO/IOC/IAEA, 1993. Reference Method No 6.
- [47] Araya, M., Olivares, M. and Pizarro, F., 2007. Copper in human health. *Int J Environ Health Res*, 1(4):608–620.
- [48] Allen, T.M., Manoli, A. and LaMont, R.L., 1982. Skeletal changes associated with copper deficiency. *Clin Orthop Relat Res*, 168:206–210.
- [49] Conlan, D., Korula, R. and Tallentire, D., 1990. Serum copper levels in elderly patients with femoral-neck fractures. *Age Ageing*, 19:212–214.
- [50] Uriu-Adams, J.Y. and Keen, C.L., 2005. Copper, oxidative stress, and human health. *Mol Aspects Med*, 26:268–298.
- [51] National Research Council (NRC), 2000. Copper in Drinking Water. Washington, DC: The National Academies Press. <https://doi.org/10.17226/9782>.
- [52] Duda-Chodak, A. and Błaszczuk, U., 2008. The impact of nickel on human health. Review paper. *J Elementol*, 13(4):685–696.
- [53] Abbaspour, N., Hurrell, R. and Kelishadi, R., 2014. Review on iron and its importance for human health. *J Res Med Sci*, 19(2):164-74.