



Synthesis Structure Elucidation and Antimicrobial Activity of Fe (II) Complex

Hardikkumar D. Chaudhary, Jwalant J. Vora and Jabali J. Vora

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

June 15, 2020

SYNTHESIS, STRUCTURE ELUCIDATION AND ANTIMICROBIAL ACTIVITY OF Fe (II) COMPLEX.

HARDIKKUMAR D. CHAUDHARY^{*}, JWALANT J.VORA

^{*}*Department of Chemistry,*

M.G. Science Institute, Navrangpura, Ahmedabad-380009, Gujarat, India

JABALI J. VORA⁺

⁺*Department of Chemistry,*

Hemchandracharya North Gujarat University, Patan - 384 265.Gujarat, India

⁺*Author for Correspondence*

E-mail: Hardikchaudhary1710@gmail.com

ABSTRACT- The conjunction of Fe ions with biologically essential and not as much of explored kynurenic acid ligand to form metal complex is an important area of modern research. These complexes were characterized by elemental analysis, molar conductance measurements, magnetic susceptibility measurements, mass spectrometry, FTIR, electronic spectral studies, TGA techniques. The stoichiometry of the complex has been found to be 1: 3 (metal: ligand). The antimicrobial activities of the complexes have been studied by screening the compounds against the bacteria *Bacillus subtilis*, *Salmonella typhi* A, *Escherichia coli* and *Staphylococcus aureus* and also results have been compared with standard drugs streptomycin and ampicillin.

Key words: Kynurenic acid, metal complexes, Antibacterial activity, Physico chemical study.

I.INTRODUCTION

Study of the metal coordination chemistry of hetero cyclic compounds is exciting, not only from an analytical point of view, but also because of the biological implications of this type of compounds.

Kynurenic acid (KYNA) was discovered in 1853 by the German chemist Justus von Liebig in dog urine, which it was apparently named after. It is produced from L-kynurenine in a reaction catalyzed by the enzyme kynurenine—oxoglutarate transaminase. Kynurenic acid is one of the endogenous products of tryptophan metabolism, formed along the kynurenine pathway. This pathway is associated between the immune and the nervous system; it also plays an immunoregulatory role during infections, pregnancies, autoimmunological processes, neoplasm growth and also after organtransplantations. Moreover, KYNA, being a selective ligand of the GPR35 receptor, is involved in the modulation of the immune response because this receptor is expressed mainly on cells connected with the immune system [1].

II. EXPERIMENTAL:

Analytical grade chemicals were used throughout the course of experimental work. Spectroscopic grade solvents were employed for recording the spectra. The compound kynurenic acid (Sigma) was used as the ligand. Fe metal carbonate used was also A.R. grade. A calculated volume of 70% HClO₄ was diluted with water to obtain 0.2M perchloric acid solution. The exact strength was firm by pH metric titration against standard 0.2M NaOH

solution. 75 ml 0.2 M perchloric acid was taken and solid Fe metal carbonate was added in it till effervescences observed (slight excess addition was done). The solution was stirred for 30 minutes and filtered and thus the metal perchlorate in aqueous solution was obtained. The formation of complex was carried out by mixing 50 mL (0.2M) metal perchlorate solution and 50 mL(0.2M) ligand in DMSO solution. The mole ratio of ligand and metal was (1:1). The reason for this ratio is lack of prior knowledge. The reaction mixture was refluxed for around 4.0 h at 90°C temperature. After 4.0 h the reaction mixture was cooled. There was no instant precipitation, then into this solution, ice water was added and straight away precipitates were obtained. The complex thus obtained was washed well with double distilled water and alcohol for elimination of unreacted metal and ligand. All the complex was dried in an oven at 40° C to 50 ° C[2]. This Fe complex was then characterized by chemical and instrumental methods to elucidate its structure.

III. RESULTS AND DISCUSSION

Table 1: RESULTS OF INFRARED SPECTRA

IR spectral band (cm ⁻¹) of KYNA ligand and Fe-KYNA complex (values mentioned in cm ⁻¹).			
KYNA	O-H Phenolic (3434), Acidic -OH (3105), Ar-CH- stretching(2967), C=N(1593), C=O Aromatic stretching(1758), bending, vibrations (748-OH out of plane), 1245, 1264 CH, CH ₂ , OH in plane, 1380- wagging and twisting).		
	Changed	New peaks	Eliminated
Fe-KYNA	O-H Phenolic(3200), Ar-CH- stretching (3085), C=N(1446), C=O Aromatic stretching (1654), bending vibrations(1446),	M-N, M-O (521,664)	Acidic -OH

Table 2: RESULTS OF PHYSICO CHEMICAL MEASUREMENTS

TLC (solvent toluene: methanol 7:3) and M.P. was taken by melting point apparatus. Metal complex formations was confirmed by TLC single spot reading. The UV – visible spectra were measured on a UV-1800 Shimadzu (Double beam) spectrophotometer.

Complex	Colour	M.W. (gm/mol)	M.P. ° C	R.F. value *	Molar Conductance (Ω) mho cm ⁻¹	% yield
Ligand (KYNA)	Light cream	189.17	269	0.8503	2.55 × 10 ⁻³	-----
Fe-KYNA	cream	629.51	279	0.8625	2.43 × 10 ⁻³	29.96

* Solvent system : (Toluene: methanol 7:3),).

Table 3: MAGNETIC AND ELECTRONIC SPECTRA.

Metal Complexes	Uv-vis spectral λ _{max} (nm)	Magn. Sus. (BM)	Number of unpaired electrons	Oxidation No.	Coordination No.	Probable shape
KYNA	346.50 291.50 258.00	-----	-----	-----	-----	-----
Fe-KYNA	360.00 345.00 296.50	6.16	4 (hs)	(II)	6	octahedral

Uv-vis = ultra violet- visible, Magn. Sus. = magnetic susceptibility, hs = high spin.

Table 4: CHN AND METAL ANALYSIS


Elemental analyses were performed with a Vario-MICRO CUBE C, H, N analyzer.

Metal Complex	C (%)		H (%)		N (%)		Metal (%) by TGA	
	Found	calculated	found	calculated	found	calculated	found	calculated
Fe-KYNA	62.21	57.46	3.85	3.35	7.18	6.70	8.57	9.41

The three columns of IR spectra results, single RF values in TLC, electronic spectra, magnetic susceptibility, elemental analyses, molar conductance, all indicated formation of complex compound in case of Fe(II). Furthermore, the over all results suggested six coordination with octahedral shape.

TGA- DSC ANALYSIS

Fig 1: Fe-KYNA



It is observed that at 150⁰C temperature in Fe-KYNA complex weight loss per mole is 2.38 gm, which indicates that no H₂O molecule of crystallization with Fe-KYNA is present and 7.95 gm weight loss occurred for one mole Fe-KYNA complex at 250⁰C temperature, which indicates that no water molecules coordinated with Fe²⁺ complex.

Table 5: Results of TGA

Compound	RT-150 °C (Water of crystallization)			150 °C – 250 °C (water of coordination)		
	% Loss	Loss of weight(gm) for 1 mole complex	water molecules	% Loss	Loss of weight(gm) for 1 mole complex	water molecules
KYNA						
Co-KYNA	0.54	2.41	0	1.77	7.91	0

RT= Room Temperature

RESULTS OF MASS SPECTROMETRY

Fig 2: [KYNA mass spectra]





Fig 3: [Fe-KYNA mass spectra]



Mass spectra:

Mass spectrum of KYNA ligand

ES⁺: 189 amu is the peak for C₁₀H₈NO₃ and 79amu is the base peak due to (C₆H₆)H⁺ and it is present because of removal of C₄HNO₃ from the hetero cyclic part.

ES⁻: 144 amu is the base peak because of removal of -COOH group from the KYNA molecule. Feeble 189 peak is observed.

Mass spectrum of Fe-KYNA

ES⁺: 157 amu is a peak due to removal of O₂ (deoxygenation) from KYNA

79 amu is the base peak due to (C₆H₆)H⁺

59 amu is a peak due to metal (Fe).

ES⁻: 248 amu is a peak because of Fe - KYNA.H⁺

188 amu is the base peak due to (KYNA - H⁺)

145 amu peak because of removal of (KYNA – COO H)

FLUORESCENCE SPECTRA

Fig 4: Combined Fluorescence spectra of metal complex and Legend





Fig.4. shows the fluorescence spectra of the ligand and the complex. The spectra are of emission type and taken in the range of 360 nm to 650 nm. The metal ion generally do not exhibit fluorescence activity under normal conditions. However the ligand KYNA indeed exhibits fluorescence activity, therefore it was considered to study the fluorescence behavior of the complex. The ligand KYNA exhibits UV-visible absorption with λ_{\max} below 400 nm but the fluorescence peaks around 405 nm and 480 nm. On co-ordination with metal ion, the fluorescence diminishes to a great extent. The probable reason seems to be due to change in the $\pi(\pi)$ bonding and lone pair electron sharing for the metal coordination.

IV.STRUCTURE

Based upon the physico chemical analyses, the structures of the Fe complex can be shown as below.

Fig 5: Fe-KYNA structure



V. ANTIMICROBIAL ACTIVITY

In vitro biological screening effects of the compounds under investigation were carried out and that was tested against the bacteria: *Salmonella typhii*, *Staphylococcus aureus*, *Escherichia coli* and *Bacillus subtilis* [3]. The well-diffusion method, using agar nutrient as the medium was used to check antimicrobial activity. In this method, generally compound was loaded into well made in agar plates and their activities were tested against different organisms streaked on the surface of agar medium [4].

The stock solution of compounds (10^{-2} M) was prepared by dissolving the compounds in DMSO. The given cultures were streaked on agar plates and wells were made with the help of cork borers. The well was filled with the test solution using a micropipette and the plate was incubated 24 h for bacteria at 35°C. During this period, the test solution diffused and the growth of the inoculated microorganisms was affected. The inhibition zone was developed, at which the concentration was noted [5].

ANTIBACTERIAL ACTIVITIES OF ANTIBIOTIC

Zone size	Antimicrobial disc used in practical
+++ 2.6 to 3.0 cm	Streptomycin (25µg/disc) for E-coli, S.typhy and S. aureus
++ 2.0 to 2.5 cm	Ampicilin (25µg/disc) for Bacillus sp.
- No zone 0.8 cm	

All antibiotics in standard condition gave +++ results.

Table 6: RESULTS OF ANTIBACTERIAL STUDIES

Culture	Well no.	Fe-KYNA	KYNA
Bacillus subtilis (Gram-possitive)	1. 25 mcg/ml	+	+
	2. 50 mcg/ml	+	+
	3. 75 mcg/ml	+	++
	4. 100 mcg/ml	+	+++
Staphylococcus aureus (Gram-possitive)	1. 25 mcg/ml	-	-
	2. 50 mcg/ml	-	-
	3. 75 mcg/ml	-	+
	4. 100 mcg/ml	++	++
Escherichiacoli (Gram negative)	1. 25 mcg/ml	-	-
	2. 50 mcg/ml	-	+
	3. 75 mcg/ml	-	-
	4. 100 mcg/ml	++	-
Salmonella typhii A (Gram negative)	1. 25 mcg/ml	-	-
	2. 50 mcg/ml	-	-
	3. 75 mcg/ml	-	-
	4. 100 mcg/ml	-	-

The KYNA has inhibitory effect on gram +ve bacteria namely *Bacillus subtilis*, *Staphylococcus* but no inhibitory effect on gram -ve cultures. The ligand and complex did not exhibit any inhibitory effect on gram negative bacteria in general. Another observation was that on coordination of KYNA, the overall antibacterial activity decreased. The antibacterial activity of ligand and complex was much less compared to the standard antibiotics selected. The ligand showed higher antibacterial activity which increased with increased concentration compared to the complex.

VI. CONCLUSION

Biological activities of Fe^{+2} as well as of the ligand, kynurenic acid, led us to prepare their complexes which showed 6 coordination number. The complexes are found to be fairly stable and these were characterized by important instrumental and routine chemical methods. Fluorescence analysis showed that on complexation, the fluorescence behaviour of the ligand was reduced but not totally eliminated. Although the antimicrobial activities the complex is found to be quite low but we earnestly are hopeful of getting encouraging results in other applications of this complex.

VII. ACKNOWLEDGEMENT

One of the authors, Jwalant vora is thankful to GUJCOST for the financial support in the form of a minor research project (GUJCOST/MRP/14-15/1460). Authors are also thankful to the principal and faculty members of M.G. Science institute, Ahmedabad. Our sincere thanks to Dr. Devang Pandya for his valuable discussions.

VIII. REFERENCES

1. J. Małaczewska¹, A. K. Siwicki¹, R. M. Wójcik¹, E. Kaczorek¹, W. A. Turski², *Cent Eur J Immunol*, V- 39(1), 2014
2. H. R. Patel, H.D. Chaudhari and J. J. Vora, *Journal of Applicable Chemistry*, 4 (6): 1774-1790, 2015
3. Irobi O N, Moo-Young M and Anderson W A 1996, *Tropical Journal of Pharmaceutical Research*, 6 (4): 809-813 December 2007
4. *Microbiology Concept and Application*, (1st edn), McGraw-Hill Inc.: New York, U.S.A; 80 – 100, 158 – 161, 370, 1993
5. Raman N, D. Raja J and Sakthivel A, *Journal of Chemical Sciences*, Volume 119, [Issue- 4](#), pp 303–310, July 2007